

EGU25
2 February 2025

Improving the CLASSIC Snow Model to Better Simulate Arctic Snowpacks


Mickaël Lalande

Postdoc at Université du Québec à Trois-Rivières (Canada) / RIVE / GLACIOLAB

ESA CCI Fellowship (SnowC²) — 01/10/2023 to 30/09/2025 (2 years)

supervised by Christophe Kinnard and Alexandre Roy


Context: Arctic Amplification



Cohen et al., (2014)

- The **Arctic** has warmed **2 to 3 times faster** than the global average (e.g., Cohen et al., [2014](#)); nearly **four times faster** than the globe since 1979 (Rantanen et al., [2022](#))
- ⇒ **melting** of **Arctic sea ice** and spring **snow cover**
- Impacts on **ecosystems** and **human activities** such as transportation, resource extraction, **water supply**, use of land and **infrastructure** among others.
- **1.035 Pg-C** (>66° N, 3m soil) - By 2100, **55 to 232 Pg C-CO₂-e** could be emitted via **permafrost degradation** (Schuur et al., [2022](#))

Snow: essential component of the climate system



Current snow models cannot simulate Arctic snowpacks!



Domine et al., (2019)




Figure 3. Comparison of measured snow density profiles at Bylot Island in May 2015 with those simulated using the detailed snow models Crocus and SNOWPACK. Crocus runs of 6 May are shown because Crocus simulates melting on 7 May, and this extra process makes comparisons irrelevant on 12 May.

Domine et al., (2018)

PHYSICAL SOLUTION

Implement the water vapor fluxes explicitly in the snowpack (→ snow mass redistribution):

- [IVORI](#) project (Marie Dumont, ERC ~2M €)
- Jafari et al., ([2020](#)): The Impact of Diffusive Water Vapor Transport on Snow Profiles in Deep and Shallow Snow Covers and on Sea Ice
- Simson et al. ([2021](#)): Elements of future snowpack modeling – Part 2: A modular and extendable Eulerian-Lagrangian numerical scheme for coupled transport, phase changes and settling processes

Arctic snowpack: solution?

PHYSICAL SOLUTION

Implement the water vapor fluxes explicitly in the snowpack (→ snow mass redistribution):

- [IVORI](#) project (Marie Dumont, ERC ~2M €)
- Jafari et al., ([2020](#)): The Impact of Diffusive Water Vapor Transport on Snow Profiles in Deep and Shallow Snow Covers and on Sea Ice
- Simson et al. ([2021](#)): Elements of future snowpack modeling – Part 2: A modular and extendable Eulerian-Lagrangian numerical scheme for coupled transport, phase changes and settling processes

PRACTICAL SOLUTION


Increase the compaction due to the wind + reduce the density of the lower layers, e.g.:

- Royer et al. ([2021](#)): Improved Simulation of Arctic Circumpolar Land Area Snow Properties and Soil Temperatures
- Lackner et al., ([2022](#)): Snow properties at the forest-tundra ecotone: predominance of water vapor fluxes even in deep, moderately cold snowpacks

Challenge: never applied worldwide and often site specific...

CLASSIC land surface model (LSM): description


Primary water, energy, CO₂, and CH₄ fluxes in CLASSIC Melton et al. (2020), Fig. 1



- **CLASSIC v1.0 LSM:** Canadian Land Surface Scheme Including Biogeochemical Cycles (Melton et al., 2020)
- → couples **CLASS 3.6.2** (Verseghy et al., 2017) and **CTEM 2.0** (Melton & Arora, 2016)
 - CLASS: physics (energy/water fluxes), etc.
 - CTEM: photosynthesis, carbon cycle, etc.
- → **used operationally** within the Canadian Earth System Model (CanESM; Swart et al., 2019) for **climate change impact assessment** (CMIP6, SnowMIP, Global Carbon Project, etc.)

CLASSIC land surface model (LSM): description

Primary water, energy, CO₂, and CH₄ flux




Snow model description (Bartlett et al., [2006](#); Brown et al., [2006](#); Langlois et al., [2014](#); Verseghy et al., [2017](#) - version 2.7 → 3.6.1):


- Single-layer snow model
- Quadratic temperature profile within the snowpack
- Snow albedo decreases and the snow density increases exponentially with time
- Fresh snow density is determined as a function of the air temperature (Pomeroy & Gray, [1995](#))
- The snow thermal conductivity is derived from the snow density (Sturm et al., [1997](#))
- Percolation and refreezing taken into account
- Interception of snowfall by vegetation is explicitly modeled (Bartlett et al., [2006](#))

CLASS: one of the best snow model

Menard et al., (2021)



CLASS → one of the **best** performing **model** in the last **SnowMIP** experiments! (SWE, SD, albedo, soil temperatures, etc.)




But performs quite **bad** for the **surface temperature**...

Main objectives / questions

- ❖ Can a **single-layer** snow model reproduce the **bulk Arctic snow** characteristic?
- ❖ Can **Arctic snowpack adaptations** be implemented **without deteriorating** model performance **elsewhere** (e.g., mid-latitude Alpine snowpacks)?
- ❖ How do those adaptations impact the **soil temperatures** and **carbon fluxes**?

Methods

Methods: 1D simulations at SnowMIP and Arctic sites



Methods: Model and simulations set-up

Forcing:

- For each site: incoming shortwave and longwave radiation, air temperature, precipitation rate (total and **solid**), air pressure, specific humidity, and wind speed
 - → linearly interpolated to the model time step (30 minutes; see [issue](#) with 1h)
 - → quality-controlled data, including correction for wind-induced solid precipitation undercatch

Initialization and boundary conditions:

- Soil properties (sand, clay, and organic matter), soil permeable depth, soil color index (SoilGrids250m), CLASS and CTEM PFTs, greenhouse gas concentration, etc.
(note: no moss and lichen, so a peat layer was added to the first soil layer (10 cm) in certain cases, e.g., at Bylot)


Spin-up:

- First spin-up 100 to 300 years (with spinfast = 10) until reaching carbon balance (looping over the full forcing files period)
- Final spin-up same duration (spinfast = 1)
- CO2 concentration fixed to the first year forcing file value


Snow model improvements

Physics improvements


#1 Soil conductivity under snow (bug)



#2 Bottom snow temperature



#3 Windless exchange coefficient



Source: Brown et al., (2006)

Arctic adaption

#4 Blowing snow sublimation losses




Credit: Les Anderson/ Unsplash

#5 Increasing snow compaction



Credit: Sawtooth Avalanche Center


#6 Snow thermal conductivity



Source: Calonne et al. (2011)

Wind effect on snow compaction: max snow density

The **snow density** increase towards a $\rho_{s,max}$ value as:
$$\rho_s(t + 1) = [\rho_s(t) - \rho_{s,max}] e^{-\frac{0.01\Delta t}{3600}} + \rho_{s,max}$$




Depends on snow depth

- Brown et al. (2006) dry snow
- Brown et al. (2006) wet snow
- Tabler et al. (1990)

Wind effect on snow compaction: max snow density

The **snow density** increase towards a $\rho_{s,max}$ value as:
$$\rho_s(t + 1) = [\rho_s(t) - \rho_{s,max}]e^{-\frac{0.01\Delta t}{3600}} + \rho_{s,max}$$



Depends on snow depth


- Brown et al. (2006) dry snow
- Brown et al. (2006) wet snow
- Tabler et al. (1990)



Problem: for typical **Arctic snowpack** (~50 cm) $\rho_{s,max}$ limited to about 200 to 250 kg m⁻³ while they usually range from 250 to 400 kg m⁻³ under strong **wind** conditions for dry snow (e.g., Domine et al., [2021](#); Royer et al., [2021a](#)).


Wind effect on snow compaction: max snow density

The **snow density** increase towards a ρ_{\max} value as:
$$\rho_s(t + 1) = [\rho_s(t) - \rho_{s,\max}] e^{-\frac{0.01\Delta t}{3600}} + \rho_{s,\max}$$




Wind effect on snow compaction: max snow density

The **snow density** increase towards a $\rho_{s,max}$ value as:
$$\rho_s(t + 1) = [\rho_s(t) - \rho_{s,max}] e^{-\frac{0.01\Delta t}{3600}} + \rho_{s,max}$$




Results


Example: Bylot example






Example: Bylot example











Example: Bylot example












Snow thermal conductivity: Sturm et al. (1997) → Calonne et al. (2011)



Overall results at all sites: RMSE







Overall results at all sites: RMSE



Conclusion

Take home message

- Improved simulated snowpack at both SnowMIP and Arctic sites!
- Snow depth biases reduced from 11.1 to 0.2 cm at the Arctic sites.
- Improved simulated snow density and temperature.
- Reduces the RMSE of the simulated soil temperatures from 5.2~°C to 3.1~°C on average at all Arctic sites.
- Simulate better soil temperature → winter carbon fluxes.
- Still uncertainties related to vegetation, soil properties, snow drifts, precip, etc. + need more snow Arctic obs sites!
- Future studies over the whole Arctic with spatial simulation + new SCF parameterizations.





MICKAËL LALANDE



SOCIAL NETWORKS



@LalandeMickael



@mickaellalande



@mickaellalande



mickaellalande.github.io

EMAIL: MICKAEL.LALANDE@UQTR.CA

Bibliography

References

- Bartlett, P. A., MacKay, M. D., & Verseghy, D. L. (2006). Modified snow algorithms in the Canadian land surface scheme : Model runs and sensitivity analysis at three boreal forest stands. *Atmosphere-Ocean*, 44(3), 207-222. <https://doi.org/10.3137/ao.440301>
- Brown, R., Bartlett, P., MacKay, M., & Verseghy, D. (2006). Evaluation of snow cover in CLASS for SnowMIP. *Atmosphere-Ocean*, 44(3), 223-238. <https://doi.org/10.3137/ao.440302>
- Calonne, N., Flin, F., Morin, S., Lesaffre, B., du Roscoat, S. R., & Geindreau, C. (2011). Numerical and experimental investigations of the effective thermal conductivity of snow. *Geophysical Research Letters*, 38(23). <https://doi.org/10.1029/2011GL049234>
- Cohen, J., Screen, J. A., Furtado, J. C., Barlow, M., Whittleston, D., Coumou, D., Francis, J., Dethloff, K., Entekhabi, D., Overland, J., & Jones, J. (2014). Recent Arctic amplification and extreme mid-latitude weather. *Nature Geoscience*, 7(9), 627-637. <https://doi.org/10.1038/ngeo2234>
- Domine, F., Belke-Brea, M., Sarrazin, D., Arnaud, L., Barrere, M., & Poirier, M. (2018). Soil moisture, wind speed and depth hoar formation in the Arctic snowpack. *Journal of Glaciology*, 64(248), 990-1002. <https://doi.org/10.1017/jog.2018.89>
- Domine, F., Picard, G., Morin, S., Barrere, M., Madore, J.-B., & Langlois, A. (2019). Major Issues in Simulating Some Arctic Snowpack Properties Using Current Detailed Snow Physics Models : Consequences for the Thermal Regime and Water Budget of Permafrost. *Journal of Advances in Modeling Earth Systems*, 11(1), 34-44. <https://doi.org/10.1029/2018MS001445>
- Domine, F., Lackner, G., Sarrazin, D., Poirier, M., & Belke-Brea, M. (2021). Meteorological, snow and soil data (2013–2019) from a herb tundra permafrost site at Bylot Island, Canadian high Arctic, for driving and testing snow and land surface models. *Earth System Science Data*, 13(9), 4331-4348. <https://doi.org/10.5194/essd-13-4331-2021>
- Jafari, M., Gouttevin, I., Couttet, M., Wever, N., Michel, A., Sharma, V., Rossmann, L., Maass, N., Nicolaus, M., & Lehning, M. (2020). The Impact of Diffusive Water Vapor Transport on Snow Profiles in Deep and Shallow Snow Covers and on Sea Ice. *Frontiers in Earth Science*, 8. <https://www.frontiersin.org/articles/10.3389/feart.2020.00249>

References

- Keenan, E., Wever, N., Dattler, M., Lenaerts, J. T. M., Medley, B., Kuipers Munneke, P., & Reijmer, C. (2021). Physics-based SNOWPACK model improves representation of near-surface Antarctic snow and firn density. *The Cryosphere*, 15(2), 1065-1085. <https://doi.org/10.5194/tc-15-1065-2021>
- Lackner, G., Domine, F., Nadeau, D. F., Lafaysse, M., & Dumont, M. (2022). Snow properties at the forest-tundra ecotone : Predominance of water vapor fluxes even in deep, moderately cold snowpacks. *The Cryosphere*, 16(8), 3357-3373. <https://doi.org/10.5194/tc-16-3357-2022>
- Langlois, A., Bergeron, J., Brown, R., Royer, A., Harvey, R., Roy, A., Wang, L., & Thériault, N. (2014). Evaluation of CLASS 2.7 and 3.5 Simulations of Snow Properties from the Canadian Regional Climate Model (CRCM4) over Québec, Canada*. *Journal of Hydrometeorology*, 15(4), 1325-1343. <https://doi.org/10.1175/JHM-D-13-055.1>
- Liston, G. E., Haehnel, R. B., Sturm, M., Hiemstra, C. A., Berezovskaya, S., & Tabler, R. D. (2007). Simulating complex snow distributions in windy environments using SnowTran-3D. *Journal of Glaciology*, 53(181), 241-256. <https://doi.org/10.3189/172756507782202865>
- Melton, J. R., & Arora, V. K. (2016). Competition between plant functional types in the Canadian Terrestrial Ecosystem Model (CTEM) v. 2.0. *Geoscientific Model Development*, 9(1), 323-361. <https://doi.org/10.5194/gmd-9-323-2016>
- Melton, J. R., Verseghy, D. L., Sospedra-Alfonso, R., & Gruber, S. (2019). Improving permafrost physics in the coupled Canadian Land Surface Scheme (v.3.6.2) and Canadian Terrestrial Ecosystem Model (v.2.1) (CLASS-CTEM). *Geoscientific Model Development*, 12(10), 4443-4467. <https://doi.org/10.5194/gmd-12-4443-2019>
- Melton, J. R., Arora, V. K., Wisernig-Cojoc, E., Seiler, C., Fortier, M., Chan, E., & Teckentrup, L. (2020). CLASSIC v1.0 : The open-source community successor to the Canadian Land Surface Scheme (CLASS) and the Canadian Terrestrial Ecosystem Model (CTEM) – Part 1 : Model framework and site-level performance. *Geoscientific Model Development*, 13(6), 2825-2850. <https://doi.org/10.5194/gmd-13-2825-2020>
- Menard, C. B., Essery, R., Krinner, G., Arduini, G., Bartlett, P., Boone, A., Brutel-Vuilmet, C., Burke, E., Cuntz, M., Dai, Y., Decharme, B., Dutra, E., Fang, X., Fierz, C., Gusev, Y., Hagemann, S., Haverd, V., Kim, H., Lafaysse, M., ... Yuan, H. (2021). Scientific and Human Errors in a Snow Model Intercomparison. *Bulletin of the American Meteorological Society*, 102(1), E61-E79. <https://doi.org/10.1175/BAMS-D-19-0329.1>

References


- Meyer, G., Humphreys, E. R., Melton, J. R., Cannon, A. J., & Lafleur, P. M. (2021). Simulating shrubs and their energy and carbon dioxide fluxes in Canada's Low Arctic with the Canadian Land Surface Scheme Including Biogeochemical Cycles (CLASSIC). *Biogeosciences*, 18(11), 3263-3283. <https://doi.org/10.5194/bg-18-3263-2021>
- Pomeroy, J., & Gray, D. (1995). Snowcover : Accumulation, Relocation, and Management. In *Bulletin of the International Society of Soil Science* (Vol. 7). <https://research-groups.usask.ca/hydrology/documents/reports/pomeroy-j.w.-and-gray-d.m.-snowcover-accumulation,-relocation-and-management-book-1995.pdf>
- Rantanen, M., Karpechko, A. Yu., Lipponen, A., Nordling, K., Hyvärinen, O., Ruosteenoja, K., Vihma, T., & Laaksonen, A. (2022). The Arctic has warmed nearly four times faster than the globe since 1979. *Communications Earth & Environment*, 3(1), 168. <https://doi.org/10.1038/s43247-022-00498-3>
- Royer, A., Picard, G., Vargel, C., Langlois, A., Gouttevin, I., & Dumont, M. (2021). Improved Simulation of Arctic Circumpolar Land Area Snow Properties and Soil Temperatures. *Frontiers in Earth Science*, 9(June), 1-19. <https://doi.org/10.3389/feart.2021.685140>
- Royer, A., Domine, F., Roy, A., Langlois, A., Marchand, N., & Davesne, G. (2021a). New northern snowpack classification linked to vegetation cover on a latitudinal mega-transect across northeastern Canada. *Écoscience*, 28(3-4), 225-242. <https://doi.org/10.1080/11956860.2021.1898775>
- Schuur, E. A. G., Abbott, B. W., Commane, R., Ernakovich, J., Euskirchen, E., Hugelius, G., Grosse, G., Jones, M., Koven, C., Leshyk, V., Lawrence, D., Lorant, M., Mauritz, M., Olefeldt, D., Natali, S., Rodenhizer, H., Salmon, V., Schädel, C., Strauss, J., ... Turetsky, M. (2022). Permafrost and Climate Change : Carbon Cycle Feedbacks From the Warming Arctic. *Annual Review of Environment and Resources*, 47(1), 343-371. <https://doi.org/10.1146/annurev-environ-012220-011847>
- Simson, A., Löwe, H., & Kowalski, J. (2021). Elements of future snowpack modeling – Part 2 : A modular and extendable Eulerian-Lagrangian numerical scheme for coupled transport, phase changes and settling processes. *The Cryosphere*, 15(12), 5423-5445. <https://doi.org/10.5194/tc-15-5423-2021>
- Sturm, M., Holmgren, J., König, M., & Morris, K. (1997). The thermal conductivity of seasonal snow. *Journal of Glaciology*, 43(143), 26-41. <https://doi.org/10.3189/S0022143000002781>

References


- Swart, N. C., Cole, J. N. S., Kharin, V. V., Lazare, M., Scinocca, J. F., Gillett, N. P., Anstey, J., Arora, V., Christian, J. R., Hanna, S., Jiao, Y., Lee, W. G., Majaess, F., Saenko, O. A., Seiler, C., Seinen, C., Shao, A., Sigmond, M., Solheim, L., ... Winter, B. (2019). The Canadian Earth System Model version 5 (CanESM5.0.3). *Geoscientific Model Development*, 12(11), 4823-4873. <https://doi.org/10.5194/gmd-12-4823-2019>
- Verseghy, D., Brown, R., & Wang, L. (2017). Evaluation of CLASS Snow Simulation over Eastern Canada. *Journal of Hydrometeorology*, 18(5), 1205-1225. <https://doi.org/10.1175/JHM-D-16-0153.1>
- Walter, B., Weigel, H., Wahl, S., & Löwe, H. (2024). Wind tunnel experiments to quantify the effect of aeolian snow transport on the surface snow microstructure. *The Cryosphere*, 18(8), 3633-3652. <https://doi.org/10.5194/tc-18-3633-2024>

Supplementary materials

Methods: SnowMIP and Arctic sites



Context: CLASSIC snow model physics (radiation)



1. **Surface energy budget:** $K_* + L_* - Q_H - Q_E - G(0) = 0$
 - a. $G(0)$ derived from the hypothesized **quadratic temperature profile** (depend only on $T(0) + \lambda_{\text{snow}}$)
 - b. + hypothesis: $G(\Delta z_s) = 0 \rightarrow T_{\text{surf}}$
2. Computation of the **snow temperature:**
 - a. Estimate bottom snow temperature



$$T_{\text{SNBOT}}(I) = \frac{Z_{\text{SNOW}}(I) * T_{\text{SNOW}}(I) + \text{DELZ}(1) * T_{\text{BAR}}(I,1)}{Z_{\text{SNOW}}(I) + \text{DELZ}(1)}$$
 - b. Compute $G(\Delta z_s)$ (same as for $G(0)$)
 - c. $\Delta T_s = [G(0) - G(\Delta z_s)] \Delta t / (C_s \Delta z_s)$

Note: In the computation of $G(\Delta z_s)$, $\lambda_{1,\text{top}}$ is considered as a harmonic average of the snow thermal conductivity and the one of the first soil layer.


Methods: CLASSIC snow model (albedo)

$$\alpha_s(t + 1) = [\alpha_s(t) - \alpha_{s,old}] e^{-\frac{0.01\Delta t}{3600}} + \alpha_{s,old}$$


| | Total albedo | Visible albedo | Near-IR albedo |
|--------------|--------------|----------------|----------------|
| Fresh snow | 0.84 | 0.95 | 0.73 |
| Old dry snow | 0.70 | 0.84 | 0.56 |
| Melting snow | 0.50 | 0.62 | 0.38 |




Physics + Arctic improvements: synthesis




Physics + Arctic improvements: synthesis



Overall results at all sites: MB




Overall results at all sites: MB



Windless transfer coefficient

Monin-Obukhov similarity theory → unable to explain turbulent energy exchanges over snow and ice surfaces under stable atmospheric conditions (turbulence does not shut down completely and is characterized by intermittent bursts). (Brown et al., 2006)

Weissfluhjoch Snow Surface Temperatures



Brown et al., (2006)

Solution → windless transfer coefficient (E_0) in the sensible heat flux:

$$Q_H = (\rho_{air} c_p C_H U + E_0) (T_s - \theta_a)$$

$E_0 = 2 \text{ W m}^{-2} \text{ K}^{-1}$ if $T_s < \theta_a$
(and $0 \text{ W m}^{-2} \text{ K}^{-1}$ otherwise)

Wind effect on snow compaction: fresh snow density

Fresh snow density in **CLASSIC**:

$$\rho_{\text{fall}} = 67.92 + 51.25 \exp(T_{\text{air}}/2.59) \quad T_{\text{air}} \leq 0^\circ\text{C} \quad (1)$$

$$\rho_{\text{fall}} = \min(200, 119.17 + 20.0 T_{\text{air}}) \quad T_{\text{air}} > 0^\circ\text{C}. \quad (2)$$

Fresh snow density in **CROCUS**:


$$\rho_{\text{new}} = \max(50, a_\rho + b_\rho(T_a - T_{\text{fus}}) + c_\rho U^{\frac{1}{2}})$$

with $a = 109 \text{ kg m}^{-3}$, $b = 6 \text{ kg m}^{-3} \text{ K}^{-1}$, and $c = 26 \text{ kg m}^{-7/2} \text{ s}^{1/2}$ → **Arctic R21 c x 2** (Royer et al., 2021)


Slight effect at the snow onset and melting but **negligible effect** on the snow depth and snow density over most of the snow season + **deterioration** at other SnowMIP sites (not shown).

→ **Snow density underestimated** of about 50 to 100 kg m^{-3}

Bylot



Arctic adaptation: Blowing snow sublimation losses



E.g. Gordon et al. (2006) → fit over multiple previous blowing snow sublimation losses parameterizations.

Total **sublimation rate**, Q_s ($\text{kg m}^{-2} \text{s}^{-1}$):


$$Q_s = A \left(\frac{T_o}{T_a} \right)^\gamma U_t \rho_a q_{si} (1 - Rh_i) (U / U_t)^B, \text{ for } U > U_t$$

and


$$U_t = U_{t*} + 0.0033(T_a - 245.88)^2$$

with $U_{t*} = 6.98 \text{ m s}^{-1}$ is the minimum threshold velocity.

Can decrease the snow depth of about ~10 cm at a few sites, but **very low impact at SnowMIP and Arctic sites**.



Arctic adaptation: Snow conductivity



CLASS snow conductivity (k_{eff}):
 → Sturm et al. (1997)

$$k_{\text{eff}} = 0.138 - 1.01\rho + 3.233\rho^2 \quad \{0.156 \leq \rho \leq 0.6\}$$

$$k_{\text{eff}} = 0.023 + 0.234\rho \quad \{\rho < 0.156\}$$

Calonne et al., (2011): "Our study, carried out on 30 snow samples spanning the full range of seasonal snow type, reveals that the effective thermal conductivity of snow is strongly correlated with snow density, and follows closely the regression curve proposed by Yen [1981]."

$$k_{\text{eff}} = 2.5 \times 10^{-6}\rho^2 - 1.23 \times 10^{-4}\rho + 0.024$$